

CO-REGULAR SPLIT DOMINATION IN GRAPHS

¹M.H.Muddebihal, ²Priyaka.H.Madarvadkar

¹Professor of Mathematics

Gulbarga University Kalaburgi -585106

Karnataka India

²Research Scholar, Department of Mathematics

Gulbarga University Kalaburgi -585106

Karnataka India

ABSTRACT: *In this paper, we introduce the new concept in domination theory. A dominating set $D \subseteq V(G)$ is a coregular split dominating set if the induced subgraph $\langle V - D \rangle$ is regular and disconnected. The minimum cardinality of such a set is called a coregular split domination number and is denoted by $\gamma_{crs}(G)$. Also we study the graph theoretic property of $\gamma_{crs}(G)$ and many bounds were obtained in terms of G and its relationship with other domination parameters were found.*

KEYWORDS: Dominating set /Split domination / Total domination/ Regular domination / Coregular split domination.

SUBJECT CLASSIFICATION NUMBER: 05C69, 05C70

1. INTRODUCTION

All graphs considered here are simple and without isolated vertices. Let $G = (V, E)$ be a graph with $|V| = P$ and $|E| = q$. We denote $\langle V - D \rangle$ to denote the subgraph induced by the set of vertices of D and $N(v)$ and $N[v]$ denote the open and closed neighborhood of a vertex v , respectively. Let $\deg(v)$ be the degree of a vertex v and as usual $\delta(G)$ the minimum degree and $\Delta(G)$ maximum degree. In general we follow the notation and terminology of Harary [2].

A vertex cover in a graph G is a set of vertices that covers all the edges of G . The vertex covering number $\alpha_o(G)$ is a minimum cardinality of a vertex cover in G . An edge cover of a graph G without isolated vertices is a set of edges of G that covers all the vertices of G . The edge covering number $\alpha_1(G)$ is a minimum cardinality of an edge cover in G .

A line graph $L(G)$ is the graph whose vertices correspond to the edges of G and two vertices in $L(G)$ are adjacent if and only if the corresponding edges in G are adjacent.

A block graph $B(G)$ is the graph whose set of vertices is the union of set of blocks of G in which two vertices are adjacent if and only if the corresponding blocks of G are adjacent.

A graph is r -regular when all its vertices have degree r , namely $\Delta(G) = \delta(G) = r$. We begin with standard definitions from domination theory.

A set $D \subseteq V$ is a dominating set of G if for every vertex $v \in V - D$, there exists a vertex $u \in D$ such that v and u are adjacent. The minimum cardinality of a dominating set in G is the domination number and denoted by $\gamma(G)$. For comprehensive work on the subject has been done in [3].

A dominating set $D \subseteq V(G)$ of a graph $G = (V, E)$ is called a connected dominating set if the induced subgraph $\langle D \rangle$ is connected. The connected domination number $\gamma_c(G)$ of G is the minimum cardinality of a connected dominating set of G see [4].

A dominating set $D \subseteq V(G)$ is a total dominating set of a graph G if the induced graph $\langle D \rangle$ does not contain an isolated vertex. The total domination number $\gamma_t(G)$ of G is the minimum cardinality of a total dominating set of G . The total domination in graph was introduced by Cockayne et al.[1] in 1980.

A dominating set $D \subseteq V(G)$ is a cototal dominating set if the induced subgraph $\langle V - D \rangle$ has no isolated vertices. The cototal domination number $\gamma_{ct}(G)$ of G is the minimum cardinality of cototal dominating set of G .

A dominating set D of G is called split dominating set if the induced subgraph $\langle V - D \rangle$ is disconnected. The split domination number is $\gamma_s(G)$ of a graph G is the minimum cardinality of a split dominating set of G .

A dominating set D of G is called strong split dominating set of G if $\langle V - D \rangle$ is totally disconnected with at least two vertices. The strong split domination number $\gamma_{ss}(G)$ of a graph G is the minimum cardinality of a strong split dominating set of G [5].

A dominating set D of G is a global dominating set if it is also dominating set of \bar{G} . A minimal cardinality of global dominating set is the global domination number and is denoted by $\gamma_g(G)$ [7].

A dominating set D of $L(G)$ is a global dominating set if it is also dominating set of $L(\bar{G})$. A minimal cardinality of D is a global domination number of $L(G)$ and denoted by $\gamma_{gl}(G)$ see[6].

2. RESULTS

We develop the following results for some standard graphs.

Theorem 1: a) For any path p_p with $p \geq 2$ vertices,

$$\gamma_{crs}(p_p) = \left\lfloor \frac{p}{2} \right\rfloor.$$

b) For any star $k_{1,p}$ with $p \geq 2$ vertices,

$$\gamma_{crs}(k_{1,p}) = 1.$$

Theorem 2: For any connected (p, q) graph G with $p \geq 3$, then

$$\gamma_{crs}(G) + \gamma(G) \leq p .$$

Proof: Let $V_1 = \{v_1, v_2, \dots, v_n\} \subseteq V(G)$ be the set of all non end vertices in G . The $V'_1 \subseteq V_1$ forms a γ -set of G . Let $V_2 = \{v_1, v_2, \dots, v_m\} \subseteq V_1$ where every $v_i \in V_2$ is adjacent to end vertices. Further $V_3 = \{v_1, v_2, \dots, v_k\} \subseteq V_1$ be the set of vertices with maximum degree. Suppose $\langle V(G) - V_2 \cup V_3 \rangle$ is disconnected and $\forall v_i \in [V(G) - \{V_2 \cup V_3\}]$ has same degree $\langle V_2 \cup V_3 \rangle$ forms a γ_{crs} -set. Otherwise there exists a set $A = \{v_1, v_2, \dots, v_k\}$ of vertices which are neighbors of some vertices in V_3 . Now $\langle V(G) - V_2 \cup V_3 \cup A \rangle$ is disconnected with isolated vertices of cardinality at least two. Then $|V_2 \cup V_3 \cup A| + |V_1| \leq V(G)$, which gives $\gamma_{crs}(G) + \gamma(G) \leq p$.

The following result gives an upper bounds for $\gamma_{crs}(G)$ in terms of γ_c and γ_t of G .

Theorem 3: For any connected (p, q) graph G with ≥ 3 , then

$$\gamma_{crs}(G) \leq \gamma_c + \gamma_t \text{ and } G \neq W_p \text{ (} P > 5\text{)}.$$

Proof: Let $V = \{v_1, v_2, \dots, v_k\}$ be the vertex set of G . Now for the graph $G \neq W_p$ with $p \geq 4$, suppose $p \leq 4$ the $\gamma_c + \gamma_t = 3 = \gamma_{crs}(G)$ and result holds. Further if $P > 5$, $|\gamma_c + \gamma_t| = 3$ and $\gamma_{crs}[W_p] = \frac{p}{2} + 1 > |\gamma_c + \gamma_t|$. Hence $G \neq W_p$ with $P > 5$. Now let $A = \{v_1, v_2, \dots, v_n\} \subseteq V(G)$ suppose for every $v \in \{V(G) - A\}$ is adjacent to at least one vertex of A . If $\langle A \rangle$ has no isolated vertices then A itself is a total dominating set of G . Otherwise let $v \in \{V(G) - A\}$ and if $\{A\} \cup \{v\}$ has no isolated vertex. Clearly $\{A\} \cup \{v\}$ is a minimal total dominating set of G . Let $A_1 = \{v_1, v_2, \dots, v_n\}$ be the set of all end vertices in G . $A_2 = \{V(G) - A_1\}$ be the set of all nonend vertices in G . Suppose there exists a minimal set of vertices such that $N[v_i] = V(G) \forall v_i \in A_2, 1 \leq i \leq n$ then A_2 forms a minimal dominating set of G . Further if $A_2 = \{V(G) - A_1\}$ has exactly one component then A_2 itself is a connected dominating set of G . Suppose A_2 has more than one component then attach the minimum set of vertices. $S' = A_2 \cup \{u, w\}$ which are in $u - w$ path, $\forall u, w \in \{V(G) - A_2\}$. Hence S' is a minimal connected dominating set of G . Further let $A_2 = \{v_1, v_2, \dots, v_i\}$ be the set of all nonend vertices suppose there exists a minimal dominating set S such that the distance between the two vertices of S is at least two clearly there exists more than one component and each component in $\langle V - S \rangle$ is regular forms γ_{crs} -set. Thus $|S| \leq |A_2| + |A|$. Hence $\gamma_{crs}(G) \leq \gamma_c + \gamma_t$.

Now the next theorem gives lower bound on the coregular split domination number of graph (G) .

Theorem 4: For any connected (p, q) graph G with $p \geq 3$, then

$$\gamma_{crs}(G) \geq \gamma_{gl}(G) - 1.$$

Proof: Let $E = \{e_1, e_2, \dots, e_n\}$ be the set of edges in G . Now consider $E_1 = \{e_1, e_2, \dots, e_k\} \subseteq E(G)$ be the set of edges with maximum edge degree and $E_2 = \{e_1, e_2, \dots, e_j\} \subseteq E(G)$ be the set of edges with minimum edge degree. Suppose $E'_1 \subseteq E_1$ and $E'_2 \subseteq E_2 \forall v \in [V(L(G)) - \{E'_1 \cup E'_2\}]$ is adjacent to at least one vertex of $\{E'_1 \cup E'_2\}$ and $\overline{\{E'_1 \cup E'_2\}}$. Since each edge of G is a vertex in $L(G)$, then $\{E'_1 \cup E'_2\}$ is a global dominating set of $L(G)$. Further let $D = \{v_1, v_2, \dots, v_n\}$ be the set of vertices in G , such that $[V(G) - N(D)]$ is regular and which gives more than one component. Then D

forms a minimal coregular split dominating set of G . Thus $|D| \geq |E'_1 \cup E'_2| - 1$ hence $\gamma_{crs}(G) \geq \gamma_{gl}(G) - 1$.

Theorem 5: For any connected (p, q) graph G with $P \geq 3$, then

$$\gamma_{crs}(G) \geq q - \alpha_1(G) + \gamma_g(G) - 1.$$

Proof: Let $A = \{v_1, v_2, v_3, \dots, v_l\}$ be set of all nonend vertices in G . Let $B_1 = \{v_1, v_2, \dots, v_m\} \subseteq A$ be a set of vertices with maximum degree. $B_2 = \{v_1, v_2, \dots, v_n\} \subseteq A$ be set of vertices with minimum degree in G . The distance between two vertices of B_1 and B_2 is at most 2. Hence $\{B_1 \cup B_2\}$ is γ -set if $[V(G) - \{B_1\} \cup \{B_2\}]$ disconnected and having vertices with same degree forms a γ -set. Let $B = \{e_1, e_2, \dots, e_n\}$ be the set of all end edges. Suppose $B' = \{e_1, e_2, \dots, e_k\} \subseteq E(G) - B$ be the set of edges such that $\text{dist}(e_i, e_j) \geq 2$ $1 \leq i \leq n$, $1 \leq j \leq k$, then $B \cup F$, where $F \subseteq B'$ be the minimal set of edges which covers all the vertices in G , such that $|B \cup F| = \alpha_1(G)$. Further let $S = \{v_1, v_2, \dots, v_p\} \subseteq V(G)$ and $S \subseteq V(\bar{G})$. If $N[S] = V(\bar{G})$. Then S is dominating set for G and (\bar{G}) . Therefore S forms a global dominating set of G . Now, we have $|B_1 \cup B_2| \leq q - |B \cup F| + |S| - 1$, which gives $\gamma_{crs}(G) \geq q - \alpha_1(G) + \gamma_g(G) - 1$.

We establish the relationship between, split domination total domination with coregular split domination number in the following theorem.

Theorem 6: For any connected (p, q) graph G with γ_{crs} is 1-regular then

$$\gamma_{crs}(G) \leq \gamma_s(G) + \gamma_t(G) - 1 \text{ and } G \neq W_p \text{ (} P > 5\text{)}.$$

Proof: Let $A_1 = \{v_1, v_2, v_3, \dots, v_n\} \subseteq V(G)$ be the set of all end vertices in G and $A'_1 = V(G) - A_1$. Suppose there exists vertex set $F \subset A'_1$ such that $D = [V(G) - F]$ is a dominating set of G . Hence $\langle D \rangle$ has more than one component with same degree than D forms a γ_{crs} -set. Suppose there exists set of vertices $C \subseteq A'_1$ where $C \cup A_1$ covers all vertices in G and if the subgraph $\langle V(G) - \{C \cup A_1\} \rangle$ does not contain any isolated vertex $C \subset A_1$ itself is a cototal dominating set of G . Otherwise if there exists a vertex $v \in [V(G) - \{C \cup A_1\}]$ with $\text{deg}(v) = 0$. Then $C \cup A_1 \cup \{v\}$ forms a minimal γ_{ct} -set of G . Further let $B' = \{v_1, v_2, \dots, v_k\} \subseteq V(G)$ be the set all nonend vertices in G . Then $B' \subseteq A'_1$ forms a minimal γ -set of G . If $\langle V - D \rangle$ is disconnected then B' forms a split dominating set of G . Hence $|D| \leq |B'| + |C| \cup A_1 \cup \{v\} - 1$ and $\gamma_{crs}(G) \leq \gamma_s(G) + \gamma_t(G) - 1$.

Theorem 7: For any non-trivial tree T with $p \geq 2$, then $\gamma_{crs}(T) = \alpha_0(T)$ if and only if γ_{crs} is zero regular.

Proof : Suppose $\gamma_{crs}(T) = \alpha_0(T)$ and γ_{csr} -set is not zero regular. Let $D = \{v_1, v_2, \dots, v_n\}$ be a dominating set of T such that the distance between two vertices of D be at most three. If $\langle V - D \rangle$ is disconnected we consider the following cases.

Case1: Assume there exists at least one edge $e \in V(T) - D$ which is a component of disconnected $\langle V(T) - D \rangle$. Then γ_{crs} is not zero regular, a contradiction.

Case2: Assume there exists a vertex $v \in \gamma_{crs}$ -set and $v \notin \alpha_0$ -set. Then there exists $N(v) = u$. Such that an edge $uv \in \{V(T) - D\}$ a contradiction.

Conversely, suppose $\gamma_{crs}(T) = \alpha_0(T)$, and $\gamma_{csr}(T)$ is zero regular. Let $D = \{v_1, v_2, \dots, v_n\}$ be a set of vertices such that the distance between two vertices of D be at most two. Hence $N(u) \cup N(v) = \varnothing, \forall u, v \in D$ and edge of T covered by the set D . Clearly $|D| = \alpha_0(T)$ since D is minimal dominating set of T and $\langle V - D \rangle$ is disconnected with $\deg(v) = 0 \forall v \in \langle V - D \rangle$. Then D is also γ_{crs} -set which is zero regular. Hence $\gamma_{crs}(T) = \alpha_0(T)$.

In the following Theorem, we establish the upper bound for $\gamma_{crs}(T)$ in terms of vertices of graph G .

Theorem 8: For any non-trivial tree T with $p \geq 2$, then $\gamma_{crs}(T) \leq p - m$. Where m is the number of end vertices in T .

Proof: Let $A = \{v_1, v_2, \dots, v_m\} \subseteq V(T)$ be the set of all end vertices in T with $|A| = m$. Let $D = \{v_1, v_2, \dots, v_n\}$ be a dominating set of T . Such that the distance between two vertices of D is at most three. If $\langle V - D \rangle$ has more than one component. Then vertices of each component have same degree and all component are also have same degree. Then D is γ_{crs} -set of a tree T . So that $|D| = p - |A|$ and gives $\gamma_{crs}(T) \leq p - m$.

Theorem 9: For any non-trivial tree T with $p \geq 2$, then $\gamma_{crs}(T) = \gamma_{ss}(T)$.

Proof: Let $H_1 = \{v_1, v_2, v_3, \dots, v_l\}$ be set of all vertices in $V(T)$. Let $H_2 = \{v_1, v_2, v_3, \dots, v_m\}$ be set of all nonend vertices adjacent to end vertices. $H_3 = \{v_1, v_2, v_3, \dots, v_n\}$ be set of all nonend vertices which are not adjacent to end vertices. Let there exists $H'_3 \subseteq H_3$ such that $D = \{H_2\} \cup \{H'_3\} \subseteq V(T)$. Where $\forall v_i \in V(T) - D$ is adjacent to at least one vertex of D . Hence D is a minimal dominating set of G . Further if $\forall v_i \in \langle V - D \rangle \deg(v_i) = 0$ with at least two vertices. Hence D is a γ_{crs} -set of G . Similarly by definition of strong split dominating set the subgraph $\langle V - D \rangle$ is a null set with at least two vertices. Hence D is also a γ_{ss} -set of G . Clearly $\gamma_{crs}(T) = \gamma_{ss}(T)$.

Further if there exists a set $E = \{e_1, e_2, \dots, e_j\}$ be edges in $\langle V - D \rangle$ and each component of $V - D$ is K_2 . Then D is a γ_{crs} -set but not γ_{ss} -set. For equality if $A = \{v_1, v_2, \dots, v_k\}$ be the set of vertices which are $N(v_m), \forall v_m \in B$ where $B = \{v_1, v_2, v_3, \dots, v_l\}$ such that $\{A\} \cup \{B\}$ forms the component as K_2 in $\langle V - D \rangle$. Then $\forall v_i \in [\{V - D\} - \{A\}]$ or $[\{V - D\} - \{B\}]$ is an isolate. Thus either $\{D\} - \{A\}$ or $\{D\} - \{B\}$ is a γ_{crs} -set and also a $\gamma_{ss}(T)$ -set of a tree. Hence $\gamma_{crs}(T) = \gamma_{ss}(T)$.

Theorem 10: For any non-trivial tree T with $p \geq 3$, then

$$\gamma_{crs}(T) + 3 \geq \left\lfloor \frac{q - \gamma_c}{2} \right\rfloor.$$

Proof: Let $V = \{v_1, v_2, \dots, v_l\}$ be vertex set of T and $E = \{e_1, e_2, \dots, e_m\}$ be edge set of T . And $A_1 = \{v_1, v_2, v_3, \dots, v_m\} \subseteq V(T)$ be set of all nonend vertices which are not adjacent to end vertices. If the distance between the two vertices of A_1 and A_2 is at most 2. Suppose there exists a set $A_2' \subseteq A_2$ hence $S = [V(T) - \{A_1 \cup A_2'\}]$ is a dominating set of T with the property that $\langle S \rangle$ is totally disconnected. Then S is a γ_{crs} -set of T . Let $H = \{A_1 \cup A_2\}$ and $\forall v_i \in V(T) - H$ is adjacent to at least one vertex of H then H is dominating set of T and $\langle H \rangle$ is connected. Hence H is γ_c -set of a tree T . Since every vertex of γ_c -set is incident with the edges of T then $(E - H)/2 \leq \{S + 3\}$, implies that $|S| + 3 \geq \left\lfloor \frac{|E| + |H|}{2} \right\rfloor$ and gives, $\gamma_{crs}(T) + 3 \geq \left\lfloor \frac{q - \gamma_c}{2} \right\rfloor$.

Next theorem gives upper bound for $\gamma_{crs}(T)$.

Theorem11: For any non-trivial tree T with $p \geq 3$, then

$$\gamma_{crs}(T) \leq \gamma_t[B(T)] + \delta(G).$$

Proof: Let $V_1 = \{v_1, v_2, v_3, \dots, v_n\}$ be the set of end vertices of $V(T)$. $V_2 = \{v_1, v_2, v_3, \dots, v_m\}$ be the set of vertices adjacent to V_1 there exists $V_3 = \{V(T) - V_1 \cup V_2\}$ then $S = \{V_2 \cup V_3\}$ is a minimal dominating set of T . Suppose there exists a $N(V_3) \cap N(V_2) = \emptyset \forall V_2, V_3 \in S$. Hence each edge of T covers by the set S and $\langle V - S \rangle$ is disconnected such that $\deg(v_i) = 0 \forall v_i \in \langle V - S \rangle$ then S is a γ_{crs} - set which is zero regular. Further let D^n be dominating set of block graph $B(T)$ of a tree T and $A_1 = V[B(T) - D^n]$ such that $D_1 \subseteq A_1$ and $\langle D_1 \cup D^n \rangle$ has no isolated vertex. Then $\{D_1 \cup D^n\}$ is γ_t - set of T . Let v be a point of minimum degree $\delta(T)$. Hence $|S| \leq |D_1 \cup D^n| + |v|$ which gives, $\gamma_{crs}(T) \leq \gamma_t[B(T)] + \delta(G)$.

In the following two lemmas we have the sharp bound attained to γ_{crs} by considering each block of G which is complete graph K_m and K_n .

Lemma 1. If G has exactly one nonend block K_n and all vertices of K_n are incident with blocks which are K_m with $m \geq n$ (or) $m < n$. Then $\gamma_{crs} = n$.

Proof: Let K_n be a nonend block of G with vertex set $D = \{v_1, v_2, \dots, v_n\}$. Suppose all vertices of K_n are incident with blocks which are K_m . We consider the following cases.

Case1: Suppose each vertex of K_n is incident with L number of blocks which are complete graphs K_m with $m \geq n$. Then D is a dominating set of G . Also the induced subgraph $\langle V(G) - D \rangle$ is disconnected and $m - 1$ regular. Hence $|D| = \gamma_{crs}(G)$, which is also equal to n . Clearly $\gamma_{crs} = n$.

Case2: Suppose each vertex of K_n is a cut vertex and incident with L number of blocks which are K_m with $m < n$. Then the induced subgraph $\langle V(G) - D \rangle$ is again disconnected and $m - 1$ regular. Since $\forall v_i \in D$ is adjacent to at least one vertex of $V(G) - D$, then D is a γ_{crs} - set of G and $|D| = n$. Clearly $\gamma_{crs} = n$.

From the above lemma we concluded that, if there exists at least one block which is either K_{m-1} or K_{m+1} in L number of blocks. Then there does not exist γ_{crs} - set.

Lemma 2: If G has exactly one cut vertex C incident with blocks which are K_n , $n \geq 2$, then $\gamma_{crs} = C$.

Proof: Suppose G has exactly one cut vertex v which is incident with m number of K_n ($n \geq 2$) blocks. Then every vertex of $\{G - V\}$ is adjacent to v . Thus $\{v\}$ is a γ - set of G and $\langle G - V \rangle$ is disconnected with m number of K_{n-1} blocks. Hence each component of $\langle G - V \rangle$ is K_{n-1} regular and $\{v\}$ is a γ_{crs} - set of G . Since v is a cut vertex then $\gamma_{crs} = C$.

Theorem12: For any graph G with C cut vertices $\gamma_{crs} = C$ if and only if G has exactly one nonend block K_n incident with complete blocks which are K_{n-c+1} .

Proof: Suppose $\gamma_{crs} = C$. Let $H = \{B_1, B_2, \dots, B_n\}$ be the set of n blocks of G . Let $A_1 = \{B_1, B_2, \dots, B_p\}$ be the end blocks in G . Such that $K_n = H - A_1$ which is nonend block of G . Let $\{v_1, v_2, \dots, v_n\} = V[K_n]$. Suppose $L_1 = \{v_1, v_2, v_3, \dots, v_i\} \subseteq V[K_n]$ be the set of cut vertices. We consider the following cases. Let D be a γ_{crs} - set of G .

Case1: Suppose $|L_1|$ cut vertices are incident with blocks which are K_{n-c} . Then L_1 is dominating set of G . But $\langle V(G) - L_1 \rangle$ is not regular. Hence $\gamma_{crs} = L_1$, contradiction.

Case 2: Suppose $\{v_1, v_2, \dots, v_n\} \in L_1$ are incident with K_{n-c+2} blocks. Then $\{L_1\}$ is a dominating set of G . Further $\langle V(G) - \{v_1, v_2, \dots, v_n\} \rangle$ is not a regular, a contradiction.

Case 3: Suppose the number of cut vertices $|L_1| > |V[K_n] - L_1|$. Then L_1 is a dominating set of G and $\langle V[G] - L_1 \rangle$ is not regular, a contradiction.

Conversely, suppose G has $\{L_1\} = C$ cut vertices and exactly one nonend block K_n incident with complete blocks K_{n-c+1} . Then $\{L_1\}$ is a dominating set of G . Further $\langle V(G) - L_1 \rangle$ is regular with more than one component. Clearly D forms a γ_{crs} -set. Hence $|D| = |L_1|$ gives $\gamma_{crs} = C$.

Theorem 13: For any graph G with C cut vertices every nonend vertex of G is adjacent with at least one end vertex then $\gamma_{crs} = C$.

Proof: For necessary condition, let $V_1 = \{v_1, v_2, v_3, \dots, v_l\} \subseteq V(G)$ be set of all end vertices in G . Let $V_2 \subseteq \{V(G) - V_1\}$ forms a γ -set of G . And let $A = \{v_1, v_2, \dots, v_m\} \subseteq V_2$ be the set of cut vertices of G . Suppose $V_3 = \{v_1, v_2, v_3, \dots, v_n\} \subseteq V_2$ be the set of nonend vertices. Then there exists at least one vertex v_i which is not adjacent to an end vertex. Since $v_j \in N(v_i)$ and $v_j \notin V_2$ and $v_i \in V_2$ then $\langle V(G) - V_2 \rangle$ is disconnected and we consider the following cases.

Case 1: Suppose G is a tree. Then $A = \{v_1, v_2, \dots, v_n\}$ be the set of all nonend vertices which are cutvertices. Suppose there exists $V'_1 \subseteq A$ which are adjacent to end vertices of T . Now assume there exists at least one vertex $v_k \in N(V'_1)$ and $v_k \notin V_2$, since v_k is a cutvertex and $\langle V(T) - V_2 \rangle$ is disconnected and regular, then $|V_2| > |V_1|$ which gives, $\gamma_{crs} \neq C$.

Case 2: Suppose G is not a tree. Then there exists at least one block which is cycle. Let v be a vertex which is not incident with an end vertex and $v \in D$ then $\langle V - V_2 \rangle$ is not regular hence D is not a γ_{crs} -set of G . Then there exists at least one vertex $u \in \{V(G) - V_2\}$ such that $\langle V(G) - \{V_2 \cup u\} \rangle$ is regular and γ_{crs} -set of G . Hence $|V_2 \cup \{u\}| > |C|$.

For sufficient conditions, let every nonend vertex of G is adjacent with at least one end vertex. Then $V_2 = \{V(G) - V_1\}$ is a dominating set of G . Also $\langle V(G) - V_2 \rangle$ is disconnected and $\deg(v_i) = 0 \forall v_i \in \{V(G) - V_1\}$. Thus V_2 is γ_{crs} -set of G . Since every vertex of V_2 is a cut vertex, then $|V_2| = |C|$. Clearly $\gamma_{crs} = C$.

REFERENCES

- [1] E. J. Cockayne, R.M Dawes, S.T Hedetniemi, Total domination in graphs, Networks 10 (1980) 211-219.
- [2] F. Harary, Graph Theory Addison, Wesley Reading Mass, 1969.
- [3] T. W. Haynes, S.T. Hedetniemi, and P.J. Slater, Fundamentals of domination in graphs, Marcel Dekker, Inc, New York (1998).
- [4] S.T. Hedetniemi and R. Laskeer, Connected domination in graphs, Graph Theory and Combinatorics, Cambridge (Academic press London, 1984) 209-217:Mr.86e:05055.
- [5] V.R Kulli, Theory of domination in graphs, Vishwa International publications (2010).
- [6] M .H. Muddebihal, Priyanka .H .Mandarvadkar, Global domination in line graphs (2019) 648-652.
- [7] E. Sampathkumar, The global domination number of graph, J. Math phy scie.23(1989) 377-385.